

Developed by the California Chlamydia Action Coalition with the assistance of staff of the STD Control Branch of the California Department of Health Services, the University of California, San Francisco and PricewaterhouseCoopers, LLP.

TABLE OF CONTENTS

	Page
Executive Summary.....	1
Background.....	2
Screening and Diagnostic Testing for Chlamydia Infection.....	3
Treatment of Chlamydia Infections	3
Endnotes	8
Resources.....	10

Figures

Treatment of Chlamydia Infections in Non-Pregnant Females and Males.....	4
Comparison of Chlamydia Testing Technologies.....	5
Chlamydia Care Path – Asymptomatic Non-Pregnant Females.....	6
Chlamydia Care Path – Uncomplicated Symptomatic Non-Pregnant Females and Males	7

EXECUTIVE SUMMARY

Sexually transmitted *Chlamydia trachomatis* is the most common communicable disease reported in California with an estimated 600,000 new infections each year. The majority of infection is asymptomatic and goes undetected. Up to 40% of women with untreated chlamydia will develop pelvic inflammatory disease (PID), the leading cause of preventable infertility in the U.S. Early identification through annual screening and appropriate treatment can significantly reduce the short- and long-term complications in women and is cost-effective

Screening

Screen all sexually active females 25 years of age and under annually for chlamydia, consistent with U.S. Centers for Disease Control and Prevention and U.S. Preventive Services Task Force guidelines, as well as HEDIS performance measurement expectations. **Use nucleic acid amplification technology tests for screening. These tests can be performed using urine specimens**, allowing chlamydia screening even if a pelvic or genital exam is not being done.

Treatment

1. Treat infected patients with either azithromycin 1 gm p.o. as a one-time dose or doxycycline 100 mg p.o. bid for 7 days. Dispense treatment on site if possible. Instruct patients to abstain from sexual intercourse for 7 days after single dose therapy or until completion of a 7-day regimen and until 7 days after all of their sexual partners are treated.
2. Evaluate, test and treat all sexual partners within the last 2 months if possible. A new California state law allows providers to prescribe treatment for partners without an examination.
3. Report infected patients to the local health department of the jurisdiction where the patient resides.

4. Test infected patients for other sexually transmitted diseases including syphilis, gonorrhea and HIV.
5. Because chlamydia reinfection is common, consider rescreening infected patients 10 weeks to 6 months after treatment.

BACKGROUND

Chlamydia trachomatis is the most common communicable disease reported in California with an estimated 600,000 new infections each year (1-3). The majority of infections are asymptomatic and go undetected. The medical consequences and costs of infection are greatest in women, who because of undetected and untreated disease, are at greatest risk of developing serious short-term and long-term complications. Up to 40% of women with untreated chlamydia will develop pelvic inflammatory disease (PID), the leading cause of preventable infertility in the U.S. (4-6). Approximately 20% of PID patients will become infertile, 6% will have ectopic pregnancies and 18% will suffer chronic pelvic pain. In men, chlamydia causes nongonococcal urethritis and acute epididymitis (6). Chlamydia may contribute substantially to the spread of heterosexually acquired HIV infection (7-9). Chlamydia is also a common cause of conjunctivitis and pneumonia in newborns born to an infected mother (10).

Risk Profile

The primary risk factors for chlamydia are age (25 years of age or under) and unprotected sex (2,6,11). Adolescent girls have the highest rates of infection, with approximately 5-10% of sexually active girls infected in California; 20 to 25-year-old women have the second highest rate ranging from 3% to 6% (12-13). Reported rates are much higher among women, largely because women are more likely to access care and be tested than men. Risk factors include new or multiple sex partners, inconsistent use of barrier contraceptives, oral contraceptive use and douching (2,6,11,14). Chlamydia has a wide socioeconomic distribution, with high rates occurring across urban, suburban and rural settings. Studies from a variety of practice settings in California, including HMOs, have consistently demonstrated high prevalence of chlamydia (5%-15%) in younger women (12,15). For example, in California in 1999, approximately 6% of girls 15-19 years of age and 4% of women 20-24

tested in a large northern California HMO were infected (12).

Screening Rationale

An estimated nine out of ten chlamydia infections in California are currently undetected and untreated because patients have no symptoms nor reasons to seek care and providers do not screen. It is estimated that 75-80% of women with chlamydia have no symptoms (3,6), and, as a result, the disease may not be diagnosed and treated until complications develop. Early identification through annual screening and subsequent treatment can significantly reduce the medical short- and long-term complications in women and has been shown to be cost-effective (16-19). A recent randomized controlled trial of chlamydia screening and treatment in an HMO demonstrated a 56% reduction in the incidence of PID in the 12 months following this intervention (19).

Screening younger women for chlamydia is recommended by the Centers for Disease Control and Prevention (CDC), the U.S. Preventive Services Task Force and a variety of professional organizations (see Resources, page 10). Recently NCQA included *Chlamydia Screening in Women** as a HEDIS measure (20). Screening for chlamydia is simple and takes very little time. Highly sensitive and specific screening tests are available (21-23). **Tests can be performed using urine specimens, allowing chlamydia screening even if a pelvic or genital exam is not being done** (24-25).

Symptomatic Infection

Most chlamydia infections are asymptomatic, so symptoms and abnormal findings are not a good predictor of infections. In addition when symptoms such as abnormal vaginal discharge, abnormal vaginal bleeding or dyspareunia are present,

* HEDIS chlamydia screening measure definition: "The percentage of women age 16 through 26 years who were identified as sexually active, who were continuously enrolled during measurement year, and who had at least one test for chlamydia during the measurement year."

they are usually mild and non-specific, and may not prompt the patient to seek care. If symptoms occur, they may appear within 2 to 4 weeks after exposure. The physical findings most commonly associated with chlamydia infection include mucopurulent cervicitis (MPC) and/or cervical friability (i.e., easily induced cervical bleeding by a swab) (6).

SCREENING AND DIAGNOSTIC TESTING FOR CHLAMYDIA INFECTIONS *

1. **Screening Criteria for Asymptomatic Sexually Active Females** (see Chlamydia Care Path for Asymptomatic Non-Pregnant Females on page 6).

Recommended: Screen all sexually active females 25 years of age and under at the first visit and annually thereafter. **Urine testing may be utilized if the patient is not scheduled for a pelvic examination.** Amplified testing [e.g. ligase chain reaction (LCR), polymerase chain reaction (PCR), strand displacement amplification (SDA), and transcription mediated amplification (TMA)] is preferred for reasons of increased sensitivity, ease of specimen collection and patient acceptability (25, 26).

2. **Diagnostic Testing for Uncomplicated Symptomatic Infection** (see Chlamydia Care Path for Uncomplicated Symptomatic Non-Pregnant Females and Males on page 7).

Recommended: Patients with symptoms or signs associated with chlamydia, including abnormal vaginal discharge, abnormal vaginal bleeding, dyspareunia, mucopurulent cervicitis (MPC) or cervical friability in females or dysuria, urethral discharge or urethritis in males, should be tested and presumptively treated for chlamydia (26).

* **Note: Screening, diagnostic testing, and treatment for pregnant females is not addressed in this clinical practice guideline.** There is a module addressing this issue in the California Chlamydia Action Coalition Quality Improvement Toolbox.

TREATMENT OF CHLAMYDIA INFECTIONS

General Principles

1. To maximize compliance with therapy, medications for chlamydia infections should be dispensed on site, if possible.
2. To minimize further transmission of infection, patients treated for chlamydia should be instructed to abstain from sexual intercourse for 7 days after single dose therapy or until completion of a 7-day regimen.
3. To minimize the risk of re-infection, patients should also be instructed to abstain from sexual intercourse until 7 days after all of their sexual partners are treated.
4. Azithromycin is probably more cost-effective in populations with poor drug compliance, little follow-up or erratic health care seeking behavior, as it provides the opportunity for single-dose, directly observed therapy.
5. Azithromycin is approved for use in persons of all ages including adolescents and children and may be particularly beneficial for use in treating adolescents (traditionally a non-compliant population). Doxycycline has the advantage of low cost and a longer history of use.
6. All sexual partners within the last two months should also be evaluated, tested and treated. Female partners, especially, should be seen and evaluated for signs and symptoms of PID. Under California law it is the duty of the attending

physician to instruct patients with STDs “in precautionary measures for preventing the spread of the disease, the seriousness of the disease and the necessity of treatment and prolonged medical supervision” (29). Additionally the attending physician is required to “endeavor to discover the source of infection, as well as any sexual or any other intimate contacts [when] the patient was in the communicable stage of the disease” and “to make an effort, through the cooperation of the patient, to bring those cases in for examination and, if necessary, treatment”(29). The Patient-delivered Partner Therapy Law enacted January 1, 2001 states “ Notwithstanding any other provision of law, a physician, nurse practitioner, certified-nurse midwife, and physician assistant who diagnoses a sexually transmitted chlamydia infection may prescribe to that patient’s sexual partner or partners without examination of that patient’s partner or partners”.

7. All patients diagnosed with chlamydia are required to be reported to the local health department of the jurisdiction where the patient resides (30).
8. Patients testing positive for chlamydia should be tested for other sexually transmitted diseases including syphilis, gonorrhea and HIV (31). In patients where other sexually transmitted diseases are initially diagnosed, a chlamydia screen should be obtained.
9. Because chlamydia reinfection is common, ranging from 10.5% (32) to 38% (33), several authors have recommended that patients be rescreened 10 weeks to 6 months after treatment (32, 34, 35).

TREATMENT OF CHLAMYDIA INFECTIONS IN NON-PREGNANT FEMALES AND MALES

Uncomplicated chlamydia infection (Asymptomatic and Symptomatic)

Recommended Regimens:

Azithromycin 1g p.o. in a single dose
or
Doxycycline 100 mg p.o. BID for 7 days

Alternative regimens: For patients allergic to Azithromycin or Doxycycline, alternatives are Ofloxacin 300 mg p.o. BID for 7 days or Erythromycin base* 500 mg p.o. QID for 7 days or Erythromycin ethylsuccinate* 800 mg p.o. QID for 7 days.

* Test of cure is recommended 3 weeks after treatment with erythromycin because of lower efficacy.

Comparison of Chlamydia Testing Technologies (21,22,25)

	NUCLEIC ACID AMPLIFICATION TECHNOLOGY*	CELL CULTURE	DIRECT FLUORESCENT ANTIBODY (DFA)	ENZYME IMMUNOASSAY (EIA)	NUCLEIC ACID PROBE
Test Type and Brand Name	<ul style="list-style-type: none"> • <i>Ligase Chain Reaction (LCR) – LCx CT Test (Abbott)</i> • <i>Polymerase Chain Reaction (PCR) – AmpliCor CT Test (Roche)</i> • <i>Transcription Mediated Amplification (TMA) – Amplified CT Assay, APTIMA (Gen-Probe)</i> • <i>Strand Displacement Amplification (SDA) - BD Probe Tec (Becton Dickinson)</i> 				<i>PACE II (Gen-Probe)</i>
Preferred Test	Yes	No	No	No	No
Collection Site	Male and female urine , endocervical and urethral swabs	Endocervical, urethral, rectal, conjunctival, pulmonary	Endocervical, urethral, rectal, conjunctival, nasopharyngeal	Endocervical, urethral, conjunctival	Endocervical, urethral, conjunctival
Sensitivity	90-95%	75-85%	70-75%	50-75%	65-75%
Specificity	98-100%	100%	95-99%	95-99%	95-99%
Test Advantages	<p>More sensitive. Non-invasive urine specimens in addition to genital swabs. For some of these tests:</p> <ul style="list-style-type: none"> • Refrigeration during transport not required. • Single specimen for chlamydia and gonorrhea. 	Recommended test for medicolegal purposes, if available.	Internally controlled for specimen adequacy. Refrigeration during transport not required.	Automated test. Refrigeration during transport not required.	Semi-automated. One swab for both chlamydia and gonorrhea. Refrigeration during transport not required.
Test Disadvantages	Contamination possible if specimen not handled properly in the clinic or lab. More costly.	Less sensitive. Longer turn around time. Technically difficult. Specimen transport, storage times and temperatures critical. Labor intensive. QA for specimen adequacy required.	Less sensitive. Technically difficult. Labor intensive.	Poor sensitivity. QA for specimen adequacy required. Confirmatory testing recommended.	Less sensitive. QA for specimen adequacy required. Confirmatory testing recommended. Negative gray zone testing recommended.

* Examples of current nucleic acid amplification tests are provided. Others are being developed.

Note: Sensitivity somewhat lower for urine compared to cervical swab specimens in nucleic acid amplification technology


Sensitivity = the proportion of patients who truly have chlamydia for whom the test is positive.

Specificity = the proportion of patients who truly do not have chlamydia for whom the test is negative.

Adapted from: USPHS Region IX Infertility Prevention Project. Clinical Guidelines. San Francisco, CA: Region IX Infertility Prevention Project Advisory Committee, 1998.

Chlamydia Care Path

ASYMPTOMATIC NON-PREGNANT FEMALES



For more information go to
<http://www.ucsf.edu/castd>

1 Azithromycin dose: 1 gram p.o., single dose.

2 Doxycycline dose: 100 mg p.o. BID for 7 days


3 Alternative Treatment Regimen: Ofloxacin 300 mg p.o. BID for 7 days or Erythromycin base 500 mg p.o. QID for 7 days or Erythromycin ethylsuccinate 800 mg p.o. QID f or 7 days.

4 Because Chlamydia reinfection is common, ranging from 10.5% (32) to 38% (33), several authors have recommended that patients be rescreened 10 weeks to 6 months after treatment (32, 34, 35)

Chlamydia Care Path

UNCOMPLICATED SYMPTOMATIC NON-PREGNANT FEMALES and MALES

For more information go to
<http://www.ucsf.edu/castd>



1 Symptoms: Females - Abnormal Vaginal Discharge, Abnormal Vaginal Bleeding, or Dyspareunia; Males - Urethral Discharge, or Dysuria.
 2 Positives include: Positive Chlamydia Test and/or Positive Physical Exam in Females - Mucopurulent Cervicitis or Cervical Friability; Positive Chlamydia Test and/or Positive Physical Exam in Males - Urethritis.
 3 Azithromycin dose: 1 gram p.o., single dose.
 4 Doxycycline dose: 100 mg p.o. BID for 7 days
 5 Alternative Treatment Regimen: Ofloxacin 300 mg p.o..BID for 7 days or Erythromycin base 500 mg p.o. QID for 7 days or Erythromycin ethylsuccinate 800 mg p.o. QID for 7 days
 6 Because Chlamydia reinfection is common, ranging from 10.5% (32) to 38% (33), several authors have recommended that patients be rescreened 10 weeks to 6 months after treatment (32, 34, 35)

ENDNOTES

1. American Social Health Association. Kaiser Family Foundation Report: Sexually Transmitted Diseases in America: How many cases and at what cost? December 1998.
2. Groseclose SL, Zaidi AA, DeLisle SJ, Levine WC, St.Louis ME. Estimated incidence and prevalence of genital *Chlamydia trachomatis* infections in the United States, 1996. *Sex Transm Dis* 1999; 26:339-344.
3. Eng TR, Butler WT Editors; Committee on Prevention and Control of Sexually Transmitted Diseases, Institute of Medicine. The Hidden Epidemic: Confronting Sexually Transmitted Diseases, National Academy Press, Washington, DC, 1997.
4. Cates W, Wasserheit JN. Genital chlamydial infections: epidemiology and reproductive sequelae. *Am J Obstet Gynecol* 1991; 164:1771-81.
5. Centers for Disease Control and Prevention. *Chlamydia trachomatis* genital infections - United States, 1996. *MMWR* 1997; 46:193-8.
6. Stamm WE. *Chlamydia trachomatis* infections of the adult. In: Holmes KK, Sparling PF, Mardh P-A, et al., eds. *Sexually Transmitted Diseases, 3rd edition*. New York, NY: McGraw Hill; 1999:407-422.
7. Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: The contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Inf.* 1999; 75:3-17.
8. Wasserheit JN. Epidemiologic synergy: interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Trans Dis* 1992; 19:61-77.
9. Royce RA, Seña A, Cates W, Cohen MS. Sexual transmission of HIV. *New Engl J Med* 1997; 336:1072-78.
10. Hammerschlag MR. Chlamydial infections in infants and children. In: Holmes KK, Sparling PF, Mardh P-A, et al., eds. *Sexually Transmitted Diseases, 3rd edition*. New York, NY: McGraw Hill; 1999:1155-1164.
11. Park BJ, Stergachis A, Scholes D, Heidrich FE, Holmes KK, Stamm WE. Contraceptive methods and the risk of *Chlamydia trachomatis* infection in young women. *Am J Epidemiol* 1995; 142:771-8.
12. California Department of Health Services, Sexually Transmitted Disease Control Branch. Sexually Transmitted Disease in California, 1999.
13. Klausner JD, McFarland W, Bolan GA, Hernandez MT, Molitor F, Lemp GF, Cahoon-Young B, Morrow S, Ruiz J. Knock-knock: A population-based survey of risk behavior, health care access, and *Chlamydia trachomatis* infection among low-income women in the San Francisco Bay Area. *J Infect Dis* 2001; 183:1087-92.
14. Scholes D, Stergachis A, Ichikawa LE, Heidrich FE, Holmes KK. Vaginal douching as a risk factor for cervical *Chlamydia trachomatis* infection. *Obstet Gynecol* 1998; 91:993-7.
15. Boyer CB, Shafer MA, Teitle E, Wibbelsman CJ, Seeberg D, Schachter J. Sexually transmitted diseases in a health maintenance organization teen clinic. *Arch Pediatr Adolesc Med* 1999; 153:838-844.
16. Marazzo JM, Celum CL, Hillis SD, Fine D, DeLisle S, Handsfield H. Performance and cost-effectiveness of selective screening criteria for *Chlamydia trachomatis* infection in women: implications for a national chlamydia control strategy. *Sex Transm Dis* 1997; 24:131-41.
17. Gene M, Mardh P-A. A cost-effectiveness analysis of screening and treatment for *Chlamydia trachomatis* infection in asymptomatic women. *Ann Intern Med.* 1996; 124:1-7.

18. Addiss DG, Vaughn ML, Ludka D, Pfister J, Davis JP. Decreased prevalence of *Chlamydia trachomatis* infection associated with a selective screening program in family planning clinics in Wisconsin. *Sex Trans Dis*. 1993; 20: 28-35.
19. Scholes D, Stergachis A, Heidrich FE, Andrilla H, Holmes KK. Prevention of pelvic inflammatory disease by screening for cervical chlamydial infection. *N Eng J Med* 1996; 334:1362-6.
20. Mangione-Smith R, McGlynn EA, Hiatt L. Screening for chlamydia in adolescents and young women. *Arch Pediatr Adolesc Med* 2000; 154:1108-13.
21. Black CM. Current methods of laboratory diagnosis of *Chlamydia trachomatis* infections. *Clin Microbiol Rev*. 1997;10:160-84.
22. Schachter J. DFA, EIA, PCR, LCR and other technologies: What tests should be used for diagnosis of chlamydia infections? *Immun Invest*. 1997; 26:157-161.
23. Stamm WE. *Chlamydia trachomatis* infections: Progress and problems. *J Infect Dis*. 1999; 179(Suppl 2):S380-3.
24. Dicker LW, Mosure DJ, Levine WC, Black CM, Berman SM. Impact of switching laboratory tests on reported trends in *Chlamydia trachomatis*. *Am J Epidemiol* 2000; 151:430-5.
25. Lindan KA. Evaluation of diagnostic tests for detecting genitourinary chlamydia infection. <http://www.ucsf.edu/castd>.
26. CDC. 1998 guidelines for the treatment of sexually transmitted diseases. *MMWR* 1998; 47 (RR-1):1-118.
27. National Committee for Quality Assurance. HEDIS 2001 Technical Specifications, Vol 2. Washington, DC: National Committee for Quality Assurance; 2000.
28. California Code of Regulations, Title 17, Chapter 4, Article 3, § 2636 (g).
29. California Code of Regulations, Title 17, Chapter 4, Article 3, § 2636 (h).
30. California Code of Regulations, Title 17, Chapter 4, Article 1, § 2500.
31. Centers for Disease Control. Sexually transmitted diseases treatment guidelines, 1989. *MMWR* 1989; 38(S-8):i-xi, 1-43.
32. Burnstein GR, Zenilman MH, Gaydos C, et al. Predictors of repeat *Chlamydia trachomatis* infections diagnosed by DNA amplification testing among inner city females. *Sex Transm Inf* 2001; 77:26-32.
33. Blythe MJ, Katz BP, Batteiger BE, Ganser JA, Jones RB. Recurrent genitourinary chlamydial infections in sexually active female adolescents. *J Pediatr* 1992; 121:487-493.
34. Whittington WLH, Kent C, Kissinger P, Oh MK, Fortenberry JD, Hillis SE, Lichtfield B, Bolan G, St. Louis ME, Farley T, Handsfield HH. Determinants of persistent and recurrent *Chlamydia trachomatis* infection in young women. *Sex Transm Dis* 2001; 28:117-23.
35. Burstein GR, Waterfield G, Joffe A, Zenilman JM, Quinn TC, Gaydos CA. Screening for gonorrhea and chlamydia by DNA amplification in adolescents attending middle school health centers. Opportunity for early intervention. *Sex Transm Dis* 1998; 25:395-402.

RESOURCES

General STD Resources

- California STD/HIV Prevention Training Center:
<http://itsa.ucsf.edu/~bolan/>
(510-883-6600)

The California STD/HIV Prevention Training Center specializes in training health care providers in the prevention, diagnosis, treatment and management of sexually transmitted diseases. Contact the center for information on course schedules and registration.
- CDC National Center for HIV/STD/TB Prevention:
<http://www.cdc.gov/nchstp/od/nchstp.html>

Home page of the National Center for Sexually Transmitted Diseases (STD), Human Immunodeficiency Virus (HIV) and Tuberculosis (TB) Prevention of the Centers for Disease Control. (CDC Voice Information System 888-CDC-FACT)
- CDC National Prevention Information Network:
<http://www.cdcnpin.org/std/start.htm>

STD Resource page of NPIN, a service of the National Center for HIV, STD and TB Prevention of the Centers for Disease Control. Links to databases, publications and other resources. (NPIN: 800-458-5231)
- The American Social Health Association:
<http://www.ashastd.org/>

Home page of ASHA, a nonprofit organization based in North Carolina dedicated to stopping STDs. Includes links to HPV/Cervical Cancer Prevention Resource Center, Herpes Resource Center and more.
- CDC STD Hotline: 800-227-8922 or 800-342-2437

Clinical Screening and Treatment Guidelines

- CDC Treatment Guidelines:
http://www.cdc.gov/nchstp/dstd/1998_STD_Guidelines/1998_guidelines_for_the_treatment.htm

The Centers for Disease Control and Prevention 1998 Guidelines for the Treatment of Sexually Transmitted Diseases.
- US Preventive Services Task Force. Screening: Chlamydial Infection. Update, 2001 Release.
<http://www.ahrq.gov/clinic/uspstf/uspstf/uspschl.htm>

Chlamydia Resources

- California Chlamydia Action Coalition:
http://www.ucsf.edu/castd/chlamydia_coalition.html

This web site includes a downloadable cost-effectiveness program to determine costs and benefits of chlamydia screening in managed care organizations. The program can be modified by chlamydia prevalence in the target population, cost of laboratory tests and cost of drugs.

Supported by the California HealthCare Foundation, based in Oakland, California. The Foundation is a non-profit philanthropic organization whose mission is to expand access to affordable, quality healthcare for underserved individuals and communities, and to promote fundamental improvements in the health status of the people of California.